

# Northumbria Research Link

Citation: O'Keefe, Jennifer M. K., Marret, Fabienne, Osterloff, Peter, Pound, Matthew and Shumilovskikh, Lyudmila (2021) Why a new volume on Non-Pollen Palynomorphs? Geological Society, London, Special Publications, 511. SP511-2021. ISSN 0305-8719

Published by: Geological Society

URL: <https://doi.org/10.1144/sp511-2021-83> <<https://doi.org/10.1144/sp511-2021-83>>

This version was downloaded from Northumbria Research Link:  
<http://nrl.northumbria.ac.uk/id/eprint/46339/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



**Northumbria  
University**  
NEWCASTLE



**UniversityLibrary**

# Why a new volume on non-pollen palynomorphs?



Jennifer M. K. O'Keefe<sup>1\*</sup>, Fabienne Marret<sup>2</sup>, Peter Osterloff<sup>3</sup>,  
Matthew J. Pound<sup>4</sup> and Lyudmila Shumilovskikh<sup>5</sup>

<sup>1</sup>Department of Physics, Earth Science, and Space Systems Engineering, Morehead State University, Morehead, Kentucky, USA

<sup>2</sup>Department of Geography and Planning, School of Environmental Sciences, University of Liverpool, Liverpool, UK

<sup>3</sup>Shell International Exploration and Production, London SE1 7NA, UK

<sup>4</sup>Department of Geography and Environmental Sciences, Northumbria University, Newcastle upon Tyne, UK

<sup>5</sup>Department of Palynology and Climate Dynamics, Georg-August-University Göttingen, Germany

✉ JMKO, 0000-0003-1793-593X; FM, 0000-0003-4244-0437;

PO, 0000-0002-4111-5336; MJP, 0000-0001-8029-9548; LS, 0000-0002-7429-3163

\*Correspondence: [j.okeefe@moreheadstate.edu](mailto:j.okeefe@moreheadstate.edu)

**Abstract:** Here we introduce the volume *Applications of Non-Pollen Palynomorphs: from Palaeoenvironmental Reconstructions to Biostratigraphy*. The study of non-pollen palynomorphs (NPPs) has a long and rich history that is interwoven with that of pollen-based studies. NPPs are among the oldest fossils on record and are instrumental in determining the origin and evolution of life, as well as studying origination and extinction events prior to the origin of pollen-producing angiosperms. This new volume on NPPs provides an up-to-date and seminal overview of the subject, linking deep-time and Quaternary study of the subject for the first time.

Why a new volume on non-pollen palynomorphs (NPPs)? Quite simply because there isn't one. Most especially not one that bridges the gaps between the use of NPPs in Quaternary and pre-Quaternary studies.

NPPs are an increasingly important part of archaeological, palaeoecological, palaeoclimatic, and biostratigraphic studies throughout the geological and archaeological record. While recognized in the fossil record for nearly two centuries, it is only in the past 75 years that marine dinoflagellate cysts have become robust biostratigraphic and palaeoecological proxies throughout the rock record, as have other marine NPPs such as Acritarchs, Prasinophytes, Chitinozoa, and Scolecodonts. Application of terrestrial NPP groups, especially in deep time, has been sporadic at best, although archaeologists, and Quaternary geologists and palaeoecologists have made significant strides in the past 30 years, thanks in a very large part to the work of Bas van Geel (1978, 1979, 1998, 2001, Fig. 1) and his students and colleagues (van Geel and van der Hammen 1978; van Geel *et al.* 1994, 1996, 2011; van Geel and Grenfell 1996; van Hove and Hendriks 1998; van Geel and Aptroot 2006, among many others). This early work catalysed Russian Quaternary

palaeoecologists (Kats *et al.* 1977). A similar effort, focused on fungi only, was seen among Canadian, Indian and American deep-time geologists (Fig. 2). However, as the focus of groups led by Jansonius (1962, 1976), Elsik (1968, 1970, 1976a, b, 1992, 1996), Elsik and Jansonius (1974), Kalgutkar (1985), Kalgutkar and Sweet (1988), Kalgutkar and McIntyre (1991), Kalgutkar and Jansonius (2000) and Saxena (1991, 2006, 2019) was largely biostratigraphic, and the majority of these scientists were not at universities, their work was not as widely adopted. Of note, both recent schools of NPP study appear to have been catalysed, at least in part, by the work of Graham (1962), who in turn had studied works stretching as far back as 1850, and Frey (1964). Recent advances in palynological processing (see Pound *et al.* 2021), changes in approach to nomenclature (see O'Keefe *et al.* 2021), especially for fossil fungi, and collaborations with mycologists (see Nuñez Otaño *et al.* 2021) and protistologists (see Andrews *et al.* 2021) have allowed the recognition of many NPPs deeper in the rock record. Indeed, some of the earliest recorded forms of life are NPPs – acid-resistant carbonaceous spheres in Archean rocks (Javeaux *et al.* 2010; Agić and Cohen 2021).


From: Marret, F., O'Keefe, J., Osterloff, P., Pound, M. and Shumilovskikh, L. (eds) *Applications of Non-Pollen Palynomorphs: from Palaeoenvironmental Reconstructions to Biostratigraphy*.

Geological Society, London, Special Publications, **511**,

<https://doi.org/10.1144/SP511-2021-83>

© 2021 The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>). Published by The Geological Society of London.

Publishing disclaimer: [www.geolsoc.org.uk/pub\\_ethics](http://www.geolsoc.org.uk/pub_ethics)




**Fig. 1.** Bas van Geel. Photograph courtesy of Encarni Montoya.

### Early history of palynology and NPP studies


The name 'non-pollen palynomorph', in many ways, is based on the assumption that pollen are most important for palynological, especially palaeoecological, studies. This is a uniquely Quaternary viewpoint. Indeed, while pollen was described by Malpighi and Grew in 1675–82 (Malpighi 1675, 1679; Grew 1682), what we now call NPPs began

to be described very soon thereafter (Fig. 3), beginning with pteridophyte and fungal spores by Tournefort in the 1690s and Geoffroy in the early 1700s (Geoffroy 1714; Stroup 1990; Bernasconi and Taiz 2006) and dinoflagellates in the mid-1700s (Baker 1753; Rochon *et al.* 2013). The first fossil pollen were described from thin-sections of coal in 1833 by Witham (1833), although they were initially described as resin vessels. The first undisputed fossil pollen were described and illustrated by line drawings in 1836 by Göppert as part of a study of fossil plants. Dinoflagellate cysts and acritarchs were described near-simultaneously by Ehrenberg (Fig. 4a; Ehrenberg 1837; Sarjeant 2002; Traverse 2007); these studies culminated with the publication of *Mikrogeologie* in 1854 (Ehrenberg 1854). Fossil microfungi were first described in 1848 (Fig. 4b; Berkeley 1848; Taylor *et al.* 2015), and their study progressed to parallel that of fossil pollen through the early 1900s. The late 1840s and early 1850s were an era of discovery; in addition to those named above, many other NPPs were identified for the first time, largely through the efforts of microscopy clubs, such as that in Clapham, UK (Sarjeant 1991); foraminiferal linings and *Botryococcus* algae were both described in 1849; prasinophytes were described in 1852; and scolecodonts in 1854 (Sarjeant 2002). In the early years, emphasis was on documenting *all* the microscopic taxa recovered in the course of a study, whether it be from a modern lake or bog or from Pleistocene peats or from Carboniferous coals. This trend continued through the earliest part of the 1900s as additional NPPs, including testate amoebae, spermatophores of copepods, and Rhabdocoelan oocytes (Rudolph 1917), were described in Quaternary bogs and peats. While changes were in the air, discovery continued, with heliozoans, *Macrobiotus* sp. eggs, and chytrids on



**Fig. 2.** Early American and Indian mycopalynologists: (a) Jan Jansonius, (b) Ramakant M. Kalgutkar, (c) William C. Elsik. Photographs (a) and (b) courtesy of AASP –The Palynological Society; used with permission; photograph (c) courtesy of Vaughn M. Bryant, Jr.

## Introduction: non-pollen palynomorphs




**Fig. 3.** A timeline of NPP discoveries. Dinoflagellate and Acritarch images from [Ehrenberg \(1837\)](#); the dinoflagellate preserves the unusual orientation chosen by Ehrenberg. Microfungi image by R. Kalgutkar in [Kalgutkar and Janssonius \(2000\)](#); used with permission of the AASP Foundation. Heliozoan image by Jablot via Wikipedia (image is in the public domain). All other palynomorph images adapted from the authors' collections.

pollen walls being described in the late 1920s ([Hesmer 1929](#)) and chitinozoans in 1931 ([Fig. 4c](#); [Eisenack 1931](#); [Sarjeant 2002](#)).

Quaternary palaeoecology began to blossom in the 1890s and early 1900s in Sweden, Denmark, Finland and Germany with the recognition that different layers of sediment from bogs preserved different quantities and assemblages of palynomorphs

([Weber 1893, 1896](#); [Lagerheim 1895](#); [Saraauw 1897](#); [Lindberg 1900](#); [Witte 1905](#); [Holst 1908](#)). With the publication of [Von Post's \(1916, 1918\)](#) papers, Quaternary palaeoecology, firmly tied to the pollen record, was born ([Edwards 2018](#)), and solidified through the work of his colleagues and students, most notably [Erdtman \(1925\)](#), [Sarjeant \(2002\)](#) and [Traverse \(2007\)](#). Despite this emphasis on



**Fig. 4.** Discoverers and describers of early NPPs: (a) Christian Gottfried Ehrenberg (1795–1876); (b) Rev. Miles Joseph Berkeley MA FLS (1803–89); (c) Alfred Eisenack (1891–1982). Painting of Ehrenberg by Eduard Radke courtesy of Wikipedia; photograph of Berkeley by Maull & Polybank, courtesy of the Wellcome Collection, Attribution 4.0 International (CC BY 4.0); photograph of Alfred Eisenack by Werner Wetzel (Tübingen) from [Gocht and Sarjeant \(1983\)](#); used with permission of *Micropaleontology*.



pollen, studies of NPPs occurring in palynology slides continued, albeit sporadically, until renewed interest in them began in the 1960s.

Palaeopalynology had begun to diverge from actuopalynology at about the same time as Quaternary palaeoecology developed, beginning with adoption of the artificial nomenclatural scheme originally proposed by Reinsch, Bennie and Kidston in the 1880s, first used by H. Potonié in the 1890s, and subsequently by palaeobotanists, especially Bartlett, in the 1920s (Bennie and Kidston 1886; Bartlett 1929*a, b*; Sarjeant 2002; see O'Keefe *et al.* 2021), and continuing with the recognition that palynomorphs could be useful in correlating coal seams beginning in 1918 (Thiessen 1918, 1920; Thiessen and Staud 1923). This realization, which came into its own in England and Germany from cross-fertilization due to visits in the mid-1920s from both Thiessen, to Sheffield and many other places (Lyons and Teichmüller 1995), and Erdtman, to various universities, including Leeds, where he worked closely with the botany and palynology group led by Burrell (Cross and Kosanke 1995; Marshall 2005). This knowledge was carried to the USA by both Erdtman himself and a young student named L.R. Wilson, who happened to be studying at the University of Leeds with Burrell immediately following Erdtman's visit (Cross and Kosanke 1995). While trained as an actuopalynologist, L.R. Wilson turned his attention to palaeopalynology beginning with a 1937 study of palynomorphs from a coal seam in Iowa (Wilson and Brokaw 1937), and collaborated with J.M. Schopf, who had himself begun to study palynomorphs in 1936, eventually producing a seminal work on Carboniferous spores in the Illinois Basin (Schopf *et al.* 1944). By 1944, however, palynostratigraphy had become the major emphasis in deep-time palynology (Wilson 1944). Interestingly, the development of palaeopalynology in Britain also began at Leeds around the same time, when A. Raistrick was a student and young researcher there. Raistrick began publishing palynological studies in the 1930s and, like Wilson, began with actuopalynological studies of peat before progressing to studies of Carboniferous spores (Raistrick and Woodhead 1930; Raistrick 1933*a, b*, 1934*a, b*, 1935, 1936, 1937, 1938, 1939; Marshall 2005). Raistrick, along with his collaborator Kathleen Blackburn, continued work on palynology ranging from Quaternary to Carboniferous studies after his move to Newcastle; a key addition in terms of NPP research was their confirmation that the Carboniferous algae noted by Thiessen many years earlier were indeed *Botryococcus*, and indistinguishable from their modern counterparts (Marshall 2005). Again, during the same period in the mid-1920s, I. Cookson was in residence in Britain, first at

Imperial College London, then at the University of Manchester (Dettmann 1993; Riding and Dettmann 2013); it is likely that she, too, was catalysed by lectures from Erdtman and Thiessen, although her interest in fossil plants and fungi was already developing. Cookson was trained as a modern botanist and mycologist in Australia, but turned her attention to palaeobotany and palaeomycology while in the UK; this collaboration produced many notable works, most importantly her seminal paper on Cenozoic fungi (1947). Near-simultaneously, in Germany, R. Potonié, beginning with his 1931 papers (Potonié 1931*a, b, c*), demonstrated the utility of palynostratigraphy and correlation in Paleogene coal-bearing sediments; these studies used the system of form-nomenclature propounded by Bartlett, which rapidly became entrenched in palaeopalynology – thus, not only were actuopalynologists and deep-time palynologists going in different directions, from the 1930s onward, they were speaking separate languages (see O'Keefe *et al.* 2021), and much of the early cross-fertilization of ideas began to wane.

Elsewhere in the world, studies of pre-Quaternary NPPs, primarily spores, acritarchs, chitinozoans, dinoflagellate cysts and prasinophytes, began in earnest in the lead-up to World War II (Sarjeant 2002). In Russia, much of this early work was led by Naumova (1939) and Liuber (1938), with an emphasis on late Paleozoic spores and pre-pollen. In India, pioneering work by Virkki (1937), a student of Birbal Sahni, on Permian floras set the stage for an explosion of palaeopalynological studies in that country. Dinoflagellate cyst studies experienced a renaissance in the 1930s, beginning with the work of Wetzel (1932, 1933*a, b*), Deflandre (1935, 1936, 1937), Lejeune (1936) and Eisenack (1931, 1935, 1936*a, b*), and Lewis (1940), and early explorations of their utility as biostratigraphic indicators by Shell Oil (Sarjeant 2002), although WWII put a hiatus on much progress. It was not until the nestor of dinoflagellate studies, W. Evitt, turned his attention to their biology and geology in the late 1950s that their study blossomed into the robust community it is today (Riding and Lucas-Clark 2016). His work catalysed fellow dinoflagellate workers Downie, Gocht, Hughes, Rossignol, Sarjeant, Vozzhennikova, Wetzel, among others (Sarjeant 2002), and led to the establishment of two major centres of fossil dinoflagellate research: (1) Stanford University in the USA and (2) the University of Sheffield in the UK. Downie's research group at Sheffield was instrumental in advancing acritarch research following WWII, as were Naumova in Moscow and Timofeyev in Leningrad, and many others in mainland Europe. Prasinophyte research did not make many advances until the post-war era, when some 14 genera were named in the period from 1952–67 (Guy-Ohlson 1996; Sarjeant 2002). It was also in

Introduction: non-pollen palynomorphs

this period that the origin of Scolecodonts was realized after Lange (1947, 1949) and Kozłowski (1956) presented articulated jaws from the Devonian of Brazil and Ordovician of Poland, respectively, and further study in Poland led to Kielan-Jaworowska’s (1966) seminal work on the preliminary phylogeny of this group. However, much of this phylogeny is now obsolete and the phylogeny and classification of scolecodonts is part and parcel of the study of fossil annelids (Parry *et al.* 2019), as is the study of clitellate cocoons, although these cocoons are of limited taxonomic value in and of themselves. Studies of Chitinozoa, too, blossomed in the post-war period, and continue to be robust biostratigraphic markers throughout their range, although their affinity remains unknown, although the consensus is that they are the remains of an extinct organism (Liang *et al.* 2019). By the late 1960s and early 1970s, study of NPPs, in both geological and Quaternary contexts was coming into its own and, through the early 2000s, has become increasingly important in palaeoecological studies.

Evolutionary history of NPPs

For much of the fossil record, it is NPPs that are dominant, and their diversification parallels the


development of multicellular life and land plants (Table 1, Fig. 5). Beginning with the simple spherical carbonaceous forms noted by Javeaux *et al.* (2010), both marine and terrestrial NPPs, including acritarchs, monolete and trilete plant spores, have been keys to understanding the oxygenation of Earth’s atmosphere (Agić and Cohen 2021), rise of multicellular life in the oceans (Agić and Cohen 2021), and the invasion of land (Wellman and Ball 2021). Indeed, Precambrian through Paleozoic biostratigraphic studies rely on NPPs, including acritarchs, chitinozoans, and precursors to dinoflagellates (Huntley *et al.* 2006; Knoll *et al.* 2007; Molyneux *et al.* 2013; Servais *et al.* 2013), as do Mesozoic and Cenozoic studies of marine sediments (Hubbard *et al.* 1994; Penaud *et al.* 2018). Thus, the evolutionary history of NPP groups is the evolutionary history of the earliest life, and a vibrant record of its diversification and preservation in the rock record.

An overview of this book

To date, no compendium addressing NPPs and their utilities from modern to ancient applications exists. This book endeavours to fill the gap by providing 12 review papers on the use and identification of NPPs. It is arranged in three sections. The first

Table 1. Geological age ranges of major groups of non-pollen palynomorphs

| Non-pollen palynomorph type             | Range in millions of years ago (Ma) | References  |
|---|-------------------------------------|---|
| Bacterial Cysts                         | 3200–Recent                         | Agić and Cohen (2021)   |
| Cyanobacteria                           | 2017–Recent                         | Hodgskiss <i>et al.</i> (2019)  |
| Achritarcha                             | 1650–Recent                         | Agić and Cohen (2021)   |
| Fungi                                   | 1230–Recent                         | Loron <i>et al.</i> (2019)  |
| Chlorophyta                             | 1000–Recent                         | Tang <i>et al.</i> (2020)   |
| Arthropoda                              | 541–Recent                          | Betts <i>et al.</i> (2014)  |
| Foraminifera (linings)                  | 540–Recent                          | Pawlowski <i>et al.</i> (2003)  |
| Scolecodonts                            | 497–Recent                          | Szaniawski (1996)   |
| Helminth eggs                           | 485–Recent                          | De Baets <i>et al.</i> (2020) preprint                                  |
| Chitinozoa                              | 480–359                             | Servais <i>et al.</i> (2013); Miller (1996)                             |
| Non-reproductive vascular plant remains | 460–Recent                          |   |
| Monolete and Trilete plant spores       | 460–Recent                          | Retallack (2020)  |
| Testate amoebae                         | 407–Recent                          | Strullu-Derrien <i>et al.</i> (2019)                                    |
| Streptophyta                            | 407–Recent                          | Head (1992), van Geel and Grenfell (1996), Wellman <i>et al.</i> (2019) |
| Freshwater sponges                      | 304–Recent                          | Schindler <i>et al.</i> (2008)  |
| Dinoflagellata                          | 247.2–Recent                        | Janouškovec <i>et al.</i> (2016)  |
| Tintinnids                              | 201.3–Recent                        | Lipps <i>et al.</i> (2013)  |
| Tardigrades                             | 145–Recent                          | Guidetti and Bertolani (2018)   |
| Loricata Euglenophyta                   | 145–Recent                          | Ascaso <i>et al.</i> (2005)   |
| Chrysophyceae                           | 112–Recent                          | Kristiansen and Škaloud (2016)  |
| Rotifers                                | 40–Recent                           | Waggoner and Poinar (1993)  |
| Rhabdocoela                             | 37.2–Recent                         | Poinar (2003) Baltic Amber  |
| Textile Fibres                          | 0.34–Recent                         | Kvavadze <i>et al.</i> (2009)   |

J. M. K. O'Keefe *et al.*

**Fig. 5.** Origin and geological age ranges of major groups of NPPs.

contains three background chapters: an overview of what organismal remains are considered NPPs (Shumilovskikh *et al.* 2021), how processing impacts the NPP spectrum obtained by the researcher (Pound *et al.* 2021) and a historical overview of nomenclature and recommendations for naming NPPs moving forward (O'Keefe *et al.* 2021). These chapters provide necessary background for current and student NPP researchers and context for interpreting what is known about NPP occurrence and utility as proxies. The second section contains an overview of the major groups of NPPs: fungi (Nuñez Otaño *et al.* 2021); freshwater remains including dinoflagellates, tintinnids, euglenids, arcellinids, rotifers thecae and eggs, flatworm egg cases, nematode eggs, and the remains of cladocerans and diptera (McCarthy *et al.* 2021); testate amoebae (Andrews *et al.* 2021); marine remains including dinoflagellates, acritarchs, tintinnids, ostracod and foraminiferal linings, copepods, and worm remains (Mudie *et al.* 2021). These chapters provide in-depth overviews of the major NPP groups in the context of their occurrence (terrestrial or marine). They are invaluable resources for understanding the intricacies of

each taxon as a proxy and interpreting their distribution in rocks and sediments. The third section provides reviews of state of the art of application of NPPs to a variety of problems: interpreting human impact on the environment (Gauthier and Jouffroy-Bapicot 2021); using coprophilous fungal spores to study megaherbivores (van Asperen *et al.* 2021); examining NPP distribution in marine settings across a major hyperthermal event (Denison 2021); tracing the origin and distribution of early land plants (Wellman and Ball 2021); and tracing the origin of early life and eukaryotes (Agić and Cohen 2021).

**Acknowledgements** Development of this book was not without its unforeseen challenges. The COVID-19 pandemic struck just as papers were being finalized for submission, significantly slowing the process as several co-authors and their families battled for their health and sanity during repeated global, regional and local shut-downs as well as a transition to primarily online course delivery and/or working remotely. We thank our many contributors, reviewers, production staff and families for their patience and support during the lengthy process.

## Introduction: non-pollen palynomorphs

**Author contributions** JMKO: conceptualization (equal), writing – original draft (lead), writing – review & editing (lead); FM: conceptualization (equal), writing – original draft (supporting), writing – review & editing (supporting); PO: conceptualization (equal), writing – review & editing (supporting); MJJP: conceptualization (equal), writing – review & editing (equal); LS: conceptualization (equal), writing – review & editing (supporting).

**Funding** This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

**Data availability** Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

## References

- Agic, H. and Cohen, P.A. 2021. Non-pollen palynomorphs in deep time: unravelling the evolution of early eukaryotes. *Geological Society, London, Special Publications*, **511**, <https://doi.org/10.1144/SP511-2020-223>
- Andrews, L.O., Payne, R.J. and Swindles, G.T. 2021. Testate amoebae as non-pollen palynomorphs in pollen slides: usefulness and application in palaeoenvironmental reconstruction. *Geological Society, London, Special Publications*, **511**, <https://doi.org/10.1144/SP511-2020-34>
- Ascaso, C., Wierzbos, J., Speranza, M., Gutiérrez, J.C., González, A.M., de los Ríos, A. and Alonso, J. 2005. Fossil protists and fungi in amber and rock substrates. *Micropalaeontology*, **51**, 59–72, <https://doi.org/10.2113/51.1.59>
- Baker, H. 1753. *Employment for the Microscope in Two Parts: Part I – An examination of salts and saline substance; Part II – An account of various animalcules*. Dodsley, London.
- Bartlett, H.H. 1929a. Fossils of the Carboniferous coal pebbles of the glacial drift at Ann Arbor. *Papers of the Michigan Academy of Science, Arts and Letters*, **9**, 11–25.
- Bartlett, H.H. 1929b. The genus *Triletes* Reinsch. *Papers of the Michigan Academy of Science, Arts and Letters*, **9**, 29–38.
- Bennie, J. and Kidston, R. 1886. On the occurrence of spores in the Carboniferous Formation of Scotland. *Proceedings of The Royal Physical Society of Edinburgh*, **9**, 82–117.
- Berkeley, M.J. 1848. On three species of mould detected by Dr. Thomas in the amber of East Prussia. *Annals and Magazine of Natural History, including Zoology, Botany, and Geology*, **2**, 380–383, <https://doi.org/10.1080/03745485809494736>
- Bernasconi, P. and Taiz, L. 2006. Claude-Joseph Geoffroy's 1711 lecture on the structure and uses of flowers. *Huntia; A Yearbook of Botanical and Horticultural Bibliography*, **13**, 1–84.
- Betts, M.J., Topper, T.P., Valentine, J.L., Skovsted, C.B., Paterson, J.R. and Brock, G.A. 2014. A new early Cambrian bradoriid (Arthropoda) assemblage from the northern Flinders Ranges, South Australia. *Gondwana Research*, **25**, 420–437, <https://doi.org/10.1016/j.jgr.2013.05.007>
- Cookson, I.C. 1947. Fossil fungi from Tertiary deposits in the Southern Hemisphere. Part I. *Proceedings of the Linnean Society of New South Wales*, **72**, 207–214.
- Cross, A.T. and Kosanke, R.M. 1995. History and development of Carboniferous palynology in North America during the early and middle twentieth century. *Geological Society of America Memoir*, **185**, 353–388.
- De Baets, K., Dentzien-Dias, P., Harrison, G.W.M., Littlewood, D.T.J. and Parry, L.A. 2020. Fossil constraints on the timescale of parasitic helminth evolution. *EcoEvoRxiv Preprints*, <https://doi.org/10.32942/osf.io/6jakv>
- Deflandre, G. 1935. Revue: Les microfossiles des silex de la craie. *Bulletin de la Société Française de Microscopie*, **4**, 116–120.
- Deflandre, G. 1936. Microfossiles des silex crétacés Pt. I: Généralités, flagellés. *Annales de Paléontologie*, **25**, 151–191.
- Deflandre, G. 1937. Microfossiles des silex crétacés Pt. II: Flagellés *incertae sedis*, hystrichosphaeridés, sarcodins, organismes divers. *Annales de Paléontologie*, **26**, 51–103.
- Denison, C.N. 2021. Stratigraphic and sedimentological aspects of the worldwide distribution of *Apectodinium* in Paleocene/Eocene Thermal Maximum deposits. *Geological Society, London, Special Publications*, **511**, <https://doi.org/10.1144/SP511-2020-46>
- Dettmann, M.E. 1993. Cookson, Isabel Clifton (1893–1973). In: Ritchie, J. (ed.) *Australian Dictionary of Biography*. National Centre of Biography, Australian National University, Canberra, **13**, 491–492.
- Edwards, K.J. 2018. Pollen, women, war and other things: reflections on the history of palynology. *Vegetation History and Archaeobotany*, **27**, 319–335, <https://doi.org/10.1007/s00334-017-0629-8>
- Ehrenberg, C.G. 1837. Die fossilen Infusorien de lebendige Dammerde. *Vorgetragen in der Akademie der Wissenschaften zu Berlin*, 1836–1837, <https://www.biodiversitylibrary.org/item/127384#page/1/mode/1up> [last accessed 15 July 2020].
- Ehrenberg, C.G. 1854. *Mikrogeologie: Das erden und felsen schaffende wirken des unsichtbar kleinen selbstständigen lebens auf der erde*. Leopold Voss, Leipzig, <https://doi.org/10.3931/e-rara-12510> [last accessed 13 July 2020].
- Eisenack, A. 1931. Neue Mikrofossilien des baltischen Silurs. *Palaeontologische Zeitschrift*, **13**, 74–118, <https://doi.org/10.1007/BF03043326>
- Eisenack, A. 1935. Mikrofossilien aus Doggergeschieben Ostpreussens. *Zeitschrift für Geschiebeforschung und Flachlandsgeologie*, **11**, 167–184.
- Eisenack, A. 1936a. Dinoflagellaten aus dem Jura. *Annales de Protistologie*, **5**, 59–63.
- Eisenack, A. 1936b. *Eodinia pachythea* n.g.n. sp., ein primitiver Dinoflagellat aus einem Kelloway-Geschiebe Ostpreussens. *Zeitschrift für Geschiebeforschung und Flachlandsgeologie*, **12**, 72–75.
- Elsik, W.C. 1968. Palynology of a Paleocene Rockdale lignite, Milam County, Texas. I. Morphology and taxonomy. *Pollen et Spores*, **10**, 1–15.



- Elsik, W.C. 1970. Palynology of a Paleocene Rockdale lignite, Milam County, Texas, III. Errata and taxonomic revisions. *Pollen et Spores*, **12**, 99–101.
- Elsik, W.C. 1976a. Fossil fungal spores. In: Weber, D.J. and Hess, W.M. (eds) *The Fungal Spore: Form and Function*. John Wiley & Sons, New York, 849–862.
- Elsik, W.C. 1976b. Fossil fungal spores and Cenozoic palynostratigraphy. *Geoscience and Man*, **15**, 115–120, <https://doi.org/10.1080/00721395.1976.9989779>
- Elsik, W.C. 1992. *The Morphology, Taxonomy, Classification and Geologic Occurrence of Fungal Palynomorphs, a Shortcourse presented under the auspices of The American Association of Stratigraphic Palynologists, Inc.* Houston, TX.
- Elsik, W.C. 1996. Chapter 10 – Fungi. In: Jansonius, J. and McGregor, D.C. (eds) *Palynology: principles and applications*. American Association of Stratigraphic Palynologists Foundation, Dallas, TX, **1**, 293–305.
- Elsik, W.C. and Jansonius, J. 1974. New genera of Paleogene fungal spores. *Canadian Journal of Botany*, **52**, 923–1150, <https://doi.org/10.1139/b74-122>
- Erdtman, G. 1925. Studies in Micro-Palaeontology. I. Evidence from the statistical study of pollen of an early Post-glacial Pine-time in North-Western Europe. 11. Moorlog from the Dogger Bank. 111. Analyses from Brittany (Finisth). IV. Peat from the Chatham Islands and the Otago District, New Zealand. *Geologiska Föreningen i Stockholm Förhandlingar*, **46**, 676–681, <https://doi.org/10.1080/11035892409454018>
- Frey, D.G. 1964. Remains of animals in Quaternary lake and bog sediments and their interpretation. *Ergebnisse der Limnologie*, **2**, 1–114.
- Gauthier, E. and Jouffroy-Bapicot, I. 2021. Detecting human impacts: non-pollen palynomorphs as proxies for human impact on the environment. *Geological Society, London, Special Publications*, **511**, <https://doi.org/10.1144/SP511-2020-54>
- Geoffroy, C.-J. 1714. Observations sur la végétation des truffes. *Histoire de l'Académie Royale des Sciences. Avec les Mémoires de Mathématique & de Physique (Paris, 4to) Année, 1711*, 23–35.
- Gocht, H. and Sarjeant, W.A.S. 1983. Pathfinder in Palynology: Alfred Eisenack (1891–1982). *Micropaleontology*, **29**, 470–477.
- Göppert, H.R. 1836. *Floribus in statu fosili comentatio*. Grassii, Bartii et Sociorum, Vratislaviae (Breslau, Poland).
- Graham, A. 1962. The role of fungal spores in palynology. *Journal of Paleontology*, **36**, 60–68.
- Grew, N. 1682. *The Anatomy of Plants; with an Idea of a Philosophical History of Plants, and several other lectures read before the Royal Society*. W. Rawlins, London
- Guidetti, R. and Bertolani, R. 2018. Paleontology and molecular dating. In: Schill, R.O. (ed.) *Water Bears: The Biology of Tardigrades*. Zoological Monographs, Springer, **2**, 131–143.
- Guy-Ohlson, D.J.E. 1996. Prasinophyceae algae. In: Jansonius, J. and McGregor, D.C. (eds) *Palynology: Principles and Applications, Vol. 1, Principles*. American Association of Stratigraphic Palynologists Foundation, Dallas, TX, 181–189.
- Head, M. 1992. Zygosporangia of the Zygnemataceae (Division Chlorophyta) and other freshwater algal spores from the uppermost Pliocene St. Erth Beds of Cornwall, southwestern England. *Micropaleontology*, **38**, 237–260, <https://doi.org/10.2307/1485790>
- Hesmer, H. 1929. Mikrofossilien in Torfen. *Paläontologische Zeitschrift*, **11**, 245–257, <https://doi.org/10.1007/BF03042729>
- Hodgskiss, M.S., Dagnaud, O.M., Frost, J.L., Halverson, G.P., Schmitz, M.D., Swanson-Hysell, N.L. and Sperling, E.A. 2019. New insights on the Orosirian carbon cycle, early cyanobacteria, and the assembly of Laurentia from the Paleoproterozoic Belcher Group. *Earth and Planetary Science Letters*, **520**, 141–152, <https://doi.org/10.1016/j.epsl.2019.05.023>
- Holst, H.O. 1908. Postglaciala tidsbestämningar. *Sveriges Geologiska Undersökning Årsbok, Series C, Afhandlingar och Uppsatser*, **2**.
- Hubbard, R.N.L.B., Boulter, M.C., and Manum, S.B. 1994. Cenozoic dinoflagellate palaeoecology elucidated, and used for marine–terrestrial biological correlation. In: Boulter, M.C. and Fisher, H.C. (eds) *Cenozoic Plants and Climates of the Arctic*. NATO ASI Series (Series I: Global Environmental Change). Springer, Berlin, **27**, [https://doi.org/10.1007/978-3-642-79378-3\\_5](https://doi.org/10.1007/978-3-642-79378-3_5)
- Huntley, J.W., Xiao, S. and Kowalewski, M. 2006. 1.3 Billion years of acritarch history: an empirical morphospace approach. *Precambrian Research*, **144**, 52–48, <https://doi.org/10.1016/j.precamres.2005.11.003>
- Jansonius, J. 1962. Palynology of Permian and Triassic sediments, Peace River area, western Canada. *Palaeontographica, Abteilung B*, **110**, 35–98.
- Jansonius, J. 1976. Paleogene fungal spores and fruiting bodies of the Canadian Arctic. *Geoscience and Man*, **15**, 129–132, <https://doi.org/10.1080/00721395.1976.9989782>
- Janouškovc, J., Gavelis, G.S. *et al.* 2016. Major transitions in dinoflagellate evolution unveiled by phylotranscriptomics. *Proceedings of the National Academy of Sciences of the United States of America*, **114**, 171–180, <https://doi.org/10.1073/pnas.1614842114>
- Javeaux, E.J., Marshall, C.P. and Bekker, A. 2010. Organic-walled microfossils in 3.2-billion-year-old shallow-marine siliciclastic deposits. *Nature*, **463**, 934–938, <https://doi.org/10.1038/nature08793>
- Kalgutkar, R.M. 1985. Fossil fungal fructifications from Bonnet Plume Formation, Yukon Territory. *Geological Survey of Canada Paper*, **85**, 259–268
- Kalgutkar, R.M. and Jansonius, J. 2000. *Synopsis of fossil fungal spores, mycelia and fructifications*. American Association of Stratigraphic Palynologists Foundation, Dallas, TX, AASP Contributions Series, **39**.
- Kalgutkar, R.M. and McIntyre, D.J. 1991. Helicosporous fungi and Early Eocene pollen, Eureka Sound Group, Axel Heiberg Island, Northwest Territories. *Canadian Journal of Earth Sciences*, **28**, 364–371, <https://doi.org/10.1139/e91-033>
- Kalgutkar, R.M. and Sweet, A.R. 1988. Morphology, taxonomy and phylogeny of the fossil fungal genus *Pesavis* from northwestern Canada. *Geological Survey of Canada, Bulletin*, **379**, 117–133.
- Kats, N.I.A., Kats, S.V. and Skobeleva, E.I. 1977. *Atlas ras-titel'nykh ostatkov v torfakh*. Nedra, Moscow.
- Kielan-Jaworowska, Z. 1966. Aparaty szczękowe wieloszczetów z ordowiku i syluru polski i porównania z

## Introduction: non-pollen palynomorphs

- formami współczesnymi. *Acta Palaeontologica Polonica*, **17**.
- Knoll, A.H., Summons, R.E., Waldbauer, J.R. and Zumberge, J.E. 2007. Chapter 8 – The geological succession of primary producers in the oceans. In: Galkowski, P.G. and Knoll, A.H. (eds) *Evolution of Primary Producers in the Sea*. Academic Press, 133–163.
- Kozłowski, R. 1956. Sur quelques appareils masticateurs des Annelides polychètes ordoviciens. *Acta Palaeontologica Polonica*, **1**, 165–205.
- Kristiansen, J. and Škaloud, P. 2016. Chrysophyta. In: Archibald, J.M., Simpson, A.G.B., Slamovits, C.H., Margulis, L., Melkonian, M., Chapman, D.J. and Corliss, J.O. (eds) *Handbook of the Protists*. Springer, Switzerland, 1–38, [https://doi.org/10.1007/978-3-319-32669-6\\_43-1](https://doi.org/10.1007/978-3-319-32669-6_43-1)
- Kvavadze, E., Bar-Yosef, O., Belfer-Cohen, A., Boaretto, E., Jakeli, N., Matskevich, Z. and Meshveliani, T. 2009. 30,000-year-old wild flax fibers. *Science (New York, NY)*, **325**, 1359, <https://doi.org/10.1126/science.1175404>
- Lagerheim, G. 1895. Uredineae Herbarii Eliae Fries. *Tromsø Museums Aarshefter*, **17**, 25–132.
- Lange, F.W. 1947. Annelidos poliquetos dos folhelhas Devonianos do Paraná Arqlivos do Museu Paranaense. *Paranaense*, **6**, 161–230.
- Lange, F.W. 1949. Polychaete annelids from the Devonian of Paraná, Brazil. *Bulletin of American Paleontology*, **33**, 5–102.
- Lejeune, M. 1936. L'étude microscopique des silex (12ième Note). *Annales de la Société Géologique Belge*, **59**, 190–197
- Lewis, H.P. 1940. The microfossils of the Upper Caradocian phosphate deposits in Montgomeryshire, North Wales. *Annals and Magazine of Natural History, Series*, **11**, 1–39
- Liang, Y., Bernardo, J., Goldman, D., Nölvak, J., Tang, P., Wang, W. and Hints, O. 2019. Morphological variation suggests that chitinozoans may be fossils of individual microorganisms rather than metazoan eggs. *Proceedings of the Royal Society B*, **286**, 20191270, <https://doi.org/10.1098/rspb.2019.1270>
- Lindberg, H. 1900. Om förekomsten i Kivinebb af subfossila växter I glaciala aflagningar. *Meddelanden af Societas pro Fauna et Flora Fennica*, **24**, 99–103
- Lipps, J.H., Stoeck, T. and Dunthorn, M. 2013. Chapter 8: Fossil Tintinnids. In: Dolan, J.R., Montagnes, D.J.S., Agatha, S., Coats, D.W. and Stoecker, D.K. (eds) *The Biology and Ecology of Tintinnid Ciliates: Models for Marine Plankton*, 1st edn. John Wiley & Sons, Oxford, 186–197.
- Liuber, A.A. 1938. Spores and pollen from the Permian of the USSR. *Problemy Sovetskoi Geologii*, **8**, 152–160 [in Russian].
- Loron, C.C., François, C., Rainbird, R.H., Turner, E.C., Borensztajn, S. and Javaux, E.J. 2019. Early fungi from the Proterozoic era in Arctic Canada. *Nature*, **570**, 232–235, <https://doi.org/10.1038/s41586-019-1217-0>
- Lyons, P.C. and Teichmüller, M. 1995. Reinhardt Thiesen (1867–1938): Pioneering coal petrologist and stratigraphic palynologist. *Geological Society of America Memoir*, **185**, 149–161, <https://doi.org/10.1130/MEM185-p149>
- Malpighi, M. 1675. *Anatome Plantarum*. Johannis Martyn, London, 1.
- Malpighi, M. 1679. *Anatome Plantarum*. Johannis Martyn, London, 2.
- Marshall, J.E.A. 2005. Arthur Raistrick: Britain's premier palynologist. *Geological Society, London, Special Publications*, **241**, 161–179, <https://doi.org/10.1144/GSL.SP.2003.207.01.13>
- McCarthy, F.M.G., Pilkington, P.M., Volik, O., Heyde, A. and Cocker, S.L. 2021. Non-pollen palynomorphs in freshwater sediments and their palaeolimnological potential and selected applications. *Geological Society, London, Special Publications*, **511**, <https://doi.org/10.1144/SP511-2020-109>
- Miller, M.A. 1996. Chapter 11. Chitinozoa. In: Jansonius, J. and McGregor, D.C. (eds) *Palynology: Principles and Applications*. American Association of Stratigraphic Palynologists Foundation, Dallas, Texas, USA, **1**, 307–336.
- Molyneux, S.G., Delabroye, A., Wicander, R. and Servais, T. 2013. Biogeography of early to mid Palaeozoic (Cambrian–Devonian) marine phytoplankton. *Geological Society, London, Memoirs*, **38**, 365–397, <https://doi.org/10.1144/M38.23>
- Mudie, P.J., Marret, F., Gurbekbe, P.R., Hartman, J.D. and Reid, P.C. 2021. Marine dinocysts, acritarchs and less well-known NPP: tintinnids, ostracod and foraminiferal linings, copepod and worm remains. *Geological Society, London, Special Publications*, **511**, <https://doi.org/10.1144/SP511-2020-55>
- Naumova, S.N. 1939. Spores and pollen of the coals of the USSR. *Transactions of the 17th International Geological Congress (1937)*, Moscow, **1**, 353–364.
- Núñez Otaño, N.B., Bianchinotti, M.V. and Saparrat, M.C.N. 2021. Palaeomycology: a modern mycological view of fungal palynomorphs. *Geological Society, London, Special Publications*, **511**, <https://doi.org/10.1144/SP511-2020-47>
- O'Keefe, J.M.K., Núñez Otaño, N.B. and Bianchinotti, M.V. 2021. Nomenclature: how do we designate NPP taxa? *Geological Society, London, Special Publications*, **511**, <https://doi.org/10.1144/SP511-2020-119>
- Parry, L.A., Eriksson, M.E. and Vinther, J. 2019. The annelid fossil record. In: Westheide, W. and Purschke, G. (eds) *Handbook of Zoology, Volume 1 Annelida Basal Groups and Pleistoannelida, Sedentaria I*. De Gruyter, Berlin, <https://doi.org/10.1515/9783110291582-003>
- Pawlowski, J., Holzmann, M. et al. 2003. The evolution of early Foraminifera. *Proceedings of the National Academy of Sciences of the United States*, **30**, 11494–11498, <https://doi.org/10.1073/pnas.2035132100>
- Penaud, A., Hardy, W. et al. 2018. Dinoflagellate fossils: Geological and biological applications. *Revue de Micropaléontologie*, **61**, 235–254, <https://doi.org/10.1016/j.revmic.2018.09.003>
- Poinar, G. 2003. A rhabdocoel turbellarian (Platyhelminthes, Typhloplanoida) in Baltic amber with a review of fossil and sub-fossil platyhelminths. *Invertebrate Biology*, **122**, 308–312, <https://doi.org/10.1111/j.1744-7410.2003.tb00095.x>
- Potonić, R. 1931a. Pollen formen aus tertiären Braunkohlen. *Jahrbuch der Preussischen Geologischen Landesanstalt zu Berlin*, **52**, 1–7.

- Potonié, R. 1931b. Zur Mikroskopie der Braunkohlen. *Tertiäre Sporen- und Blütenstaubformen Braunkohle*, **30**, 554–556.
- Potonié, R. 1931c. Pollen formen der miocänen Braunkohle. *Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin, Jahrg.*, **1931**, 24–28.
- Pound, M.J., O'Keefe, J.M.K. and Marret, F. 2021. An overview of techniques applied to the extraction of non-pollen palynomorphs, their known taphonomic issues and recommendations to maximize recovery. *Geological Society, London, Special Publications*, **511**, <https://doi.org/10.1144/SP511-2020-40>
- Raistrick, A. 1933a. The microspores of coal and their use in correlation. *Geological Magazine*, **70**, 479, <https://doi.org/10.1017/S0016756800096564>
- Raistrick, A. 1933b. The microspores of coal and their use in correlation. *Report of the British Association for the Advancement of Science for 1933*.
- Raistrick, A. 1934a. The correlation of coal-seams by microspore-content. Part I. – The seams of Northumberland. *Transactions of the Institution of Mining Engineers*, **88**, 142–153, 259–264.
- Raistrick, A. 1934b. The correlation of coal seams. *Armstrong College Mining Society Journal*, **10**, 14–22.
- Raistrick, A. 1935. The microspore analysis of coal. *The Naturalist*, **942**, 145–150.
- Raistrick, A. 1936. Use of microspores in the correlation of coal seams (Trencherbone and Busty Seams). *Report of the British Association for the Advancement of Science for 1936*, 354.
- Raistrick, A. 1937. The microspores of coal and their use in correlation, etc. *Compte rendu du deuxième Congrès pour l'avancement des études de Stratigraphie Carbonifère – Heerlen 1935*. Van Aelst, Maastricht, 909–917.
- Raistrick, A. 1938. The microspore content of some Lower Carboniferous coals. *Transactions of the Leeds Geological Association*, **5**, 221–226.
- Raistrick, A. 1939. The correlation of coal-seams by microspore-content. Part II. – The Trencherbone Seam, Lancashire, and the Busty Seams, Durham. *Transactions of the Institution of Mining Engineers*, **97**, 425–437; **98**, 95–99, 171–175.
- Raistrick, A. and Woodhead, T.W. 1930. Plant remains in post-glacial gravels near Leeds. *The Naturalist*, **877**, 39–44.
- Retallack, G.J. 2020. Ordovician land plants and fungi from Douglas Dam, Tennessee. *The Palaeobotanist*, **63**, 1–33.
- Riding, J.B. and Dettmann, M.E. 2013. The first Australian palynologist: Isabel Clifton Cookson (1893–1973) and her scientific work. *Alcheringa: An Australasian Journal of Palaeontology*, **38**, 97–129, <https://doi.org/10.1080/03115518.2013.828252>
- Riding, J.B. and Lucas-Clark, J. 2016. The life and scientific work of William R. Evitt (1923–2009). *Palynology*, **40**, 2–131, <https://doi.org/10.1080/01916122.2016.1147792>
- Rochon, A., Harland, R. and De Vernal, A. 2013. Dinoflagellates and their cysts: key foci for future research. In: Lewis, J.M., Marret, F. and Bradley, L. (eds) *Biological and Geological Perspectives of Dinoflagellates*. The Micropaleontological Society Special Publications, 89–95.
- Rudolph, K. 1917. Untersuchungen über den Aufbau Böhmischer Moore. I. Aufbau und Entwicklungsgeschichte Südböhmischer Hochmoore. *Abhandlungen der Zoologisch-Botanischen Gesellschaft in Wien*, **9**, 1–116.
- Saraauw, G.F.L. 1897. Cromer-skovlaget i Frihavnen og traellevningerne i de ravførende sandlad ved København. *Meddelelser fra Dansk Geologisk Forening*, **4**, 17–44.
- Sarjeant, W.A.S. 1991. Henry Hopley White (1790–1877) and the early researches on Chalk 'Xanthidia' (marine palynomorphs) by Clapham microscopists. *Journal of Micropalaeontology*, **10**, 83–93, <https://doi.org/10.1144/jm.10.1.83>
- Sarjeant, W.A.S. 2002. 'As chimney-sweepers, come to dust': a history of palynology to 1970. In: Oldroyd, D.R. (ed.) *The Earth Inside and Out: Some Major Contributions to Geology in the Twentieth Century*. Geological Society, London, Special Publications, **192**, 273–327, <https://doi.org/10.1144/GSL.SP.2002.192.01.15>
- Saxena, R.K. 1991. *A Catalogue of Fossil Plants from India - Part 5B. Tertiary Fungi*. Birbal Sahni Institute of Palaeobotany, Lucknow.
- Saxena, R.K. 2006. *A Catalogue of Tertiary Fungi from India. Diamond Jubilee Special Publication*. Birbal Sahni Institute of Palaeobotany, Lucknow.
- Saxena, R.K. 2019. Validation of Names of Fossil Fungi from Tertiary Sediments of India. *Novon: A Journal for Botanical Nomenclature*, **22**, 223–226, <https://doi.org/10.3417/2009095>
- Schindler, T., Wuttke, M. and Poschmann, M. 2008. Oldest record of freshwater sponges (Porifera: Spongillina) – spiculite finds in the Permo-Carboniferous of Europe. *Paläontologische Zeitschrift*, **82**, 373–384, <https://doi.org/10.1007/BF03184428>
- Schopf, J.M., Wilson, L.R. and Bentall, R. 1944. An annotated synopsis of Paleozoic fossil spores and the definition of generic groups. *Illinois Geological Survey Report of Investigations*, **91**.
- Servais, T., Achab, A. and Asselin, E. 2013. Eighty years of chitinozoan research: from Alfred Eisenack to Florentin Parin. *Review off Palaeobotany and Palynology*, **198**, 2–13, <https://doi.org/10.1016/j.revpalbo.2013.08.001>
- Shumilovskikh, L., O'Keefe, J.M.K. and Marret, F. 2021. An overview of the taxonomic groups of non-pollen palynomorphs. *Geological Society, London, Special Publications*, **511**, <https://doi.org/10.1144/SP511-2020-65>
- Stroup, A. 1990. *A Company of Scientists: Botany, Patronage, and Community at the Seventeenth-Century Parisian Royal Academy of Sciences*. University of California Press, Berkeley, <http://ark.cdlib.org/ark:/13030/ft587006gh/> [last accessed 09 July 2020].
- Strullu-Derrien, C., Kenrick, P., Goral, T. and Knoll, A.H. 2019. Testate amoebae in the 407-million-year-old Rhynie Chert. *Current Biology*, **29**, 461–467, <https://doi.org/10.1016/j.cub.2018.12.009>
- Szaniawski, H. 1996. Chapter 12: Scolecodonts. In: Jansoni, J. and McGregor, D.C. (eds) *Palynology: Principles and Applications. Volume 1: Principles*. American Association of Stratigraphic Palynologists Foundation, Dallas, 337–354.
- Tang, Q., Pang, K., Yuan, X. and Xiao, S. 2020. A one-1431 billion-year-old multicellular chlorophyte.

# Introduction: non-pollen palynomorphs

- Nature Ecology & Evolution*, **4**, 543–549, <https://doi.org/10.1038/s41559-020-1122-9>
- Taylor, T.N., Krings, M. and Taylor, E.L. 2015. *Fossil Fungi*. Academic Press, Elsevier, Amsterdam.
- Thiessen, R. 1918. The determination of the stratigraphic position of coal seams by means of their spore-exines [abs.]. *Science, New Series*, **47**, 469.
- Thiessen, R. 1920. The correlation of coal seams by means of spore-exines [abs.]. *Science, New Series*, **51**, 522.
- Thiessen, R. and Staud, J.N. 1923. Correlation of coal beds in the Monongahela Formation of Ohio, Pennsylvania, and West Virginia. *Carnegie Institute of Technology, Coal Mining Investigations Bulletin*, **9**.
- Traverse, A. 2007. *Paleopalynology*, 2nd edn. Topics in Geobiology. Springer, Dordrecht, The Netherlands, **28**.
- van Asperen, E.N., Perrotti, A. and Baker, A. 2021. Coprophilous fungal spores: non-pollen palynomorphs for the study of past megaherbivores. *Geological Society, London, Special Publications*, **511**, <https://doi.org/10.1144/SP511-2020-41>
- van Geel, B. 1978. A palaeoecological study of Holocene peat bog sections in Germany and the Netherlands, based on the analysis of pollen, spores and macro- and microscopic remains of fungi, algae, cormophytes, and animals. *Review of Palaeobotany and Palynology*, **25**, 1–120, [https://doi.org/10.1016/0034-6667\(78\)90040-4](https://doi.org/10.1016/0034-6667(78)90040-4)
- van Geel, B. 1979. Preliminary report on the history of Zygnemataceae and the use of their spores as ecological markers. *Proceedings of the IV International Palynological Conference, Lucknow (1976–1977)*, 467–469.
- van Geel, B. 1998. Are the resting eggs of the rotifer *Hexarthra mira* (Hudson 1871) the modern analogs of *Schizosporis reticulatus* Cookson and Dettmann 1959? *Palynology*, **22**, 83–87, <https://doi.org/10.1080/01916122.1998.9989504>
- van Geel, B. 2001. Non-pollen palynomorphs. In: Smol, J.P., Birks, H.J.B. and Last, W.M. (eds) *Tracking environmental change using lake sediments. Volume 3: Terrestrial, Algal, and Siliceous indicators*. Kluwer, Dordrecht, 1–17.
- van Geel, B. and Aptroot, A. 2006. Fossil ascomycetes in Quaternary deposits. *Nova Hedwigia*, **82**, 313–329, <https://doi.org/10.1127/0029-5035/2006/0082-0313>
- van Geel, B. and Grenfell, H.R. 1996. Spores of Zygnemataceae. In: Jansonius, J. and McGregor, D.C. (eds) *Palynology: principles and applications. Volume 1: Principles*. American Association of Stratigraphic Palynologists Foundation, Dallas, 173–179.
- van Geel, B. and van der Hammen, T. 1978. Zygnemataceae in Quaternary Colombian sediments. *Review of Palaeobotany and Palynology*, **25**, 377–392, [https://doi.org/10.1016/0034-6667\(78\)90021-0](https://doi.org/10.1016/0034-6667(78)90021-0)
- van Geel, B., Mur, L.R., Ralska-Jasiewiczowa, M. and Goslar, T. 1994. Fossil akinetes of Aphanizomenon and Anabaena as indicators for medieval phosphate-eutrophication of Lake Gościąg (Central Poland). *Review of Palaeobotany and Palynology*, **83**, 97–105, [https://doi.org/10.1016/0034-6667\(94\)90061-2](https://doi.org/10.1016/0034-6667(94)90061-2)
- van Geel, B., Odgaard, B.V. and Ralska-Jasiewiczowa, M. 1996. Cyanobacteria as indicators of phosphate-eutrophication of lakes and pools in the past. *Pact*, **50**, 399–415.
- van Geel, B., Gelorini, V. et al. 2011. Diversity and ecology of tropical African fungal spores from a 25 000-year palaeo-environmental record in southeastern Kenya. *Review of Palaeobotany and Palynology*, **164**, 174–190, <https://doi.org/10.1016/j.revpalbo.2011.01.002>
- van Hoeve, M.L. and Hendrikse, M. (eds) 1998. *A study of non-pollen objects in pollen slides. The types as described by Dr. Bas van Geel and colleagues*. Laboratory of Palynology and Palaeobotany, Utrecht.
- Virkki, C. 1937. On the occurrence of winged spores in the Lower Gondwana rocks of India and Australia. *Proceedings of the Indian Academy of Sciences, Section B*, **6**, 428–431.
- Von Post, L. 1916. Einige Südschwedischen Quellmoore. *Bulletin of the Geological Institute of Uppsala University*, **15**, 219–278.
- Von Post, L. 1918. Skogsträdpollen i sydsvenska torvmosselagerföljder. Förhandlingar ved de 16. Skandinavia Naturforskermøte, **1916**, 433–465.
- Waggoner, B.M. and Poinar, G.O. 1993. Fossil habrotrichid rotifers in Dominican amber. *Experientia*, **49**, 354–357, <https://doi.org/10.1007/BF01923421>
- Weber, C.A. 1893. Ober die diluviale Flora von Fahrenberg in Holstein. *Botanisches Jahrbuch*, **18**, 43
- Weber, C.A. 1896. Zur Kritik interglacialer Pflanzenablägerungen. *Abhandlungen herausgegeben vom Naturwissenschaftlichen Verein zu Bremen*, **13**, 413–468.
- Wellman, C.H. and Ball, A.C. 2021. Early land plant phyto-debris. *Geological Society, London, Special Publications*, **511**, <https://doi.org/10.1144/SP511-2020-36>
- Wellman, C.H., Graham, L.E. and Lewis, L.A. 2019. Filamentous green algae from the Early Devonian Rhynie chert. *PalZ*, **93**, 387–393, <https://doi.org/10.1007/s12542-019-00456-z>
- Wetzel, O. 1932. Die Typen der Baltischen Geschiebe-feuersteine, beurteilt nach ihrem Gehalt an Mikrofossilien. *Zeitschrift für Geschiebeforschung*, **8**, 129–146.
- Wetzel, O. 1933a. Die in organischer Substanz erhaltenen Mikrofossilien des Baltischen Kreide-Feuersteins, Teil I. *Palaeontographica, Abteilung A*, **77**, 141–188.
- Wetzel, O. 1933b. Die in organischer Substanz erhaltenen Mikrofossilien des Baltischen Kreide-Feuersteins, Teil II. *Palaeontographica, Abteilung A*, **78**, 1–110.
- Wilson, L.R. 1944. Spores and pollen as microfossils. *Botanical Review*, **10**, 499–523, <https://doi.org/10.1007/BF02861126>
- Wilson, L.R. and Brokaw, A.L. 1937. Plant microfossils of an Iowa coal deposit. *Proceedings of the Iowa Academy of Science*, **44**, 127–130.
- Witham, H.T.M. 1833. *The Internal Structure of Fossil Vegetables Found in the Carboniferous and Oolitic Deposits of Great Britain, Described and Illustrated*. Black, Edinburgh; Longman, Rees, Orme, Brown, Green & Longman, London.
- Witte, H. 1905. *Stratiotes aloides* L. funnen i Sveriges postglaciala avlagringar. *Geologiska Föreningen i Stockholm Förhandlingar*, **27**, 432, <https://doi.org/10.1080/11035890509445502>